

Random Solitons Realization of Quantum Mechanics and Stochastic Qubits

Yu.P. Rybakov¹, T.F. Kamalov²

¹ Theoretical Physics Department People's Friendship University of Russia

Mikluho-Maklay str., 6, Moscow, 117198, Russia

e-mail: yrybakov@sci.pfu.edu.ru; yrybakov@mtu-net.ru

² Physics Department Moscow State Opened University

P. Korchagin str., 22, Moscow, 107996, Russia

e-mail: ykamalov@rambler.ru

Abstract

Stochastic realization of the wave function in quantum mechanics, with the inclusion of soliton representation of extended particles, is discussed. The concept of Stochastic Qubits is used for quantum computing modeling.

1 Introduction. Wave—Particle Dualism and Solitons

As a first motivation for introducing stochastic representation of the wave function let us consider the de Broglie plane wave

$$\psi = Ae^{-\beta kx} = Ae^{-\beta\omega t + \beta(\mathbf{k}\mathbf{r})} \quad (1)$$

for a free particle with the energy ω , momentum $\mathbf{k}$, and mass m , when the relativistic relation

$$k^2 = \omega^2 - \mathbf{k}^2 = m^2 \quad (2)$$

holds (in natural units $\hbar = c = 1$).

Suppose, following L. de Broglie [1] and A. Einstein [2], that the structure of the particle is described by a regular bounded function $u(t, \mathbf{r})$, which is supposed to satisfy some nonlinear equation with the Klein — Gordon linear part. Let $l_0 = 1/m$ be the characteristic size of the soliton solution $u(t, \mathbf{r})$ moving with the velocity $\mathbf{v} = \mathbf{k}/\omega$.

Now it is worth-while to underline the remarkable fact behind this research [3], namely, the possibility to represent the de Broglie wave (1) as the sum of solitons located at nodes of a cubic lattice with the spacing $a \gg l_0$:

$$Ae^{-\beta kx} = \sum_{\mathbf{d}} u(t, \mathbf{r} + \mathbf{d}), \quad (3)$$

where $\mathbf{d}$ marks the positions of lattice nodes. To show the validity of (3) one can take into account the asymptotic behavior of the soliton in its tail region:

$$u(x) = \int d^4k e^{-\beta kx} g(k) \delta(k^2 - m^2) \quad (4)$$

and then use the well-known formula

$$\sum_{\mathbf{d}} e^{i\mathbf{b}(\mathbf{k}\mathbf{d})} = \left(\frac{2\pi}{a}\right)^3 \delta(\mathbf{k}), \quad (5)$$

implying that

$$A = \left(\frac{2\pi}{a}\right)^3 \frac{g(m)}{2m}.$$

The formula (3) gives a simple illustration of the wave — particle dualism, showing that the de Broglie wave characterizes the assemblage of particles — solitons.

2 Random Hilbert space

The next step to get the stochastic representation of quantum mechanics can be performed if one admits that the locations of solitons' centers are not regular nodes of the cubic lattice but some randomly chosen points. To realize this prescription, suppose that a field ϕ describes n particles — solitons and has the form

$$\phi(t, \mathbf{r}) = \sum_{k=1}^n \phi^{(k)}(t, \mathbf{r}), \quad (6)$$

where

$$\text{supp } \phi^{(k)} \cap \text{supp } \phi^{(k')} = 0, \quad k \neq k',$$

and the same for the conjugate momenta

$$\pi(t, \mathbf{r}) = \partial\mathcal{L}/\partial\phi_t = \sum_{k=1}^n \pi^{(k)}(t, \mathbf{r}), \quad \phi_t = \partial\phi/\partial t.$$

Let us define the auxiliary functions

$$\varphi^{(k)}(t, \mathbf{r}) = \frac{1}{\sqrt{2}}(\nu_k \phi^{(k)} + i\pi^{(k)}/\nu_k) \quad (7)$$

with the constants ν_k satisfying the normalization condition

$$\hbar = \int d^3x |\varphi^{(k)}|^2. \quad (8)$$

Now we define the analog of the wave function in the configurational space $\mathbb{R}^{3n} \ni \mathbf{x} = \{\mathbf{r}_1, \dots, \mathbf{r}_n\}$ as

$$\Psi_N(t, \mathbf{r}_1, \dots, \mathbf{r}_n) = (\hbar^n N)^{-1/2} \sum_{j=1}^N \prod_{k=1}^n \varphi_j^{(k)}(t, \mathbf{r}_k), \quad (9)$$

where $N \gg 1$ stands for the number of trials (observations) and $\varphi_j^{(k)}$ is the one-particle function (7) for the j - th trial. It can be shown [3, 4] that the quantity

$$\rho_N = \frac{1}{(\Delta V)^n} \int_{(\Delta V)^n \subset \mathbb{R}^{3n}} d^{3n}x |\Psi_N|^2,$$

where ΔV is the elementary volume which is supposed to be much greater than the proper volume of the particle $l_0^3 = v_0 \ll \Delta V$, plays the role of coordinate probability density.

If we choose the classical observable A with the generator $\hat{M}_A$, one can represent it in the form

$$A_j = \int d^3x \pi_j \hat{M}_A \phi_j = \sum_{k=1}^n \int d^3x \varphi_j^{*(k)} \hat{M}_A^{(k)} \varphi_j^{(k)},$$

for the j - th trial. The corresponding mean value is

$$\mathbb{E}(A) = \frac{1}{N} \sum_{j=1}^N A_j = \frac{1}{N} \sum_{j=1}^N \sum_{k=1}^n \int d^3x \varphi_j^{*(k)} \hat{M}_A^{(k)} \varphi_j^{(k)} = \int d^{3n}x \Psi_N^* \hat{A} \Psi_N + O\left(\frac{v_0}{\Delta V}\right), \quad (10)$$

where the hermitian operator $\hat{A}$ reads

$$\hat{A} = \sum_{k=1}^n \hbar \hat{M}_A^{(k)}. \quad (11)$$

Thus, upto the terms of the order $v_0/\Delta V \ll 1$, we obtain the standard quantum mechanical rule (10) for the calculation of mean values [4, 5].

It is interesting to underline that the solitonian scheme in question contains also the well-known spin — statistics correlation [6]. Namely, if $\varphi_j^{(k)}$ is transformed under the rotation by irreducible representation $D^{(J)}$ of $SO(3)$, with the weight J , then the transposition of two identical extended particles is equivalent to the relative 2π rotation of $\varphi_j^{(k)}$, that gives the multiplication factor $(-1)^{2J}$ in Ψ_N .

It can be also proved that Ψ_N upto the terms of order $v_0/\Delta V$ satisfies the standard Schrödinger equation [6]. To verify the fact that solitons can really possess wave properties, the *gedanken* diffraction experiment with individual electrons — solitons was realized. Solitons with some velocity were dropped into a rectilinear slit, cut in the impermeable screen, and the transverse momentum was calculated which they gained while passing the slit, the width of which significantly exceeded the size of the soliton. As a result, the picture of distribution of the centers of scattered solitons was restored on the registration screen, by considering their initial distribution to be uniform over the transverse coordinate. It was clarified that though the center of each soliton fell into a definite place of the registration screen (depending on the initial soliton profile and the point of crossing the plane of the slit by the center), the statistical picture in many ways was similar to the well-known

diffraction distribution in optics, i.e. Fresnel's picture at short distances from the slit and Fraunhofer's one at large distances [7, 8].

Fulfillment of the quantum mechanics correspondence principle for the Einstein — de Broglie's soliton model was discussed in the works [4–6]. In these papers it was shown that in the framework of the soliton model all quantum postulates were regained at the limit of point particles so that from the physical fields one can build the amplitude of probability and the average can be calculated as a scalar product in the Hilbert space by introducing the corresponding quantum operators (11) for observables. The fundamental role of the gravitational field in the de Broglie — Einstein solitonian scheme was discussed in [6, 9]. The soliton model of the hydrogen atom was developed in [10, 11].

As a result we obtain the stochastic realization (9) of the wave function Ψ_N which can be considered as an element of the random Hilbert space $\mathcal{H}_{\text{rand}}$ with the scalar product

$$(\psi_1, \psi_2) = \mathbb{E}(\psi_1^* \psi_2), \quad (12)$$

with $\mathbb{E}$ standing for the expectation value. As a rude simplification one can admit that the averaging in (12) is taken over random characteristics of particles — solitons, such as their positions, velocities, phases, and so on. It is important to underline once more that the correspondence with the standard quantum mechanics is retained only in the point — particle limit ($\Delta v \gg v_0$) for $N \rightarrow \infty$. To show this [4, 5] one can apply the central limit theorem stating that for $N \rightarrow \infty$ $\Psi_N(t, \mathbf{x})$ behaves as the Gaussian random field with the variance

$$\sigma^2 = \rho(t, \mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^{3n}, \quad (13)$$

where $\rho(t, \mathbf{x})$ stands for the probability density (partition function) of solitons' centers in $\mathbb{R}^{3n}$.

Random Hilbert spaces being widely exploited in mathematical statistics [12], for quantum applications they were first used by N. Wiener in [13, 14]. To illustrate the line of Wiener's argument, we recall the general scheme of introducing various representations in quantum mechanics.

Let $|\psi\rangle$ be a state vector in the Hilbert space $\mathcal{H}$ and $\hat{A}$ be a self-conjugate operator with the spectrum $\sigma(\hat{A})$. Then the a — representation is given by the wave function

$$\psi(a) = \langle a | \psi \rangle,$$

where

$$\hat{A}|a\rangle = a|a\rangle, \quad a \in \sigma(\hat{A}).$$

In particular, the famous Schrödinger coordinate q — representation is given by the wave function

$$\psi(q) = \langle q | \psi \rangle = \sum_n \langle q | n \rangle \langle n | \psi \rangle, \quad (14)$$

with $|n\rangle$ being some complete set of state vectors in $\mathcal{H}$.

Wiener considered the real Brownian process $x(s, \alpha)$ in the interval $[0, 1] \ni s$, where $\alpha \in [0, 1]$ is the generalized number of the Brownian trajectory and the correlator reads

$$\int_0^1 d\alpha x(s, \alpha) x(s', \alpha) = \min(s, s'). \quad (15)$$

To obtain the quantum mechanical description, Wiener defined the complex Brownian process

$$z(s|\alpha, \beta) = \frac{1}{\sqrt{2}} [x(s, \alpha) + \mathfrak{B} y(s, \beta)]; \alpha, \beta \in [0, 1], \quad (16)$$

and using the natural mapping $\mathbb{R}^3 \rightarrow [0, 1]$, for the particle in $\mathbb{R}^3$, constructed the stochastic representation of the wave function along similar lines as in (14):

$$\langle \alpha, \beta | \psi \rangle = \int_{s \in [0, 1]} dz(s|\alpha, \beta) \psi(s), \quad (17)$$

with the obvious unitarity property

$$\int_0^1 ds |\psi(s)|^2 = \iint_{[0, 1]^2} d\alpha d\beta |\langle \alpha, \beta | \psi \rangle|^2$$

stemming from (15).

3 Stochastic qubits and solitons

Now we intend to explain how stochastic qubits (quantum bits) could be introduced in the solitonian scheme. To this end one should define the random phase Φ_j for the j -th trial in our system of n solitons — particles. Let $\varphi^{(k)}(\mathbf{r})$ denote the standard (etalon) profile for the k -th soliton. The most probable position $\mathbf{d}_j^{(k)}(t)$ of the k -th soliton's center in j -th trial can be found from the following variational problem:

$$\left| \int d^3x \varphi_j^{*(k)}(t, \mathbf{r}) \varphi^{(k)}(\mathbf{r} - \mathbf{d}_j^{(k)}) \right| \rightarrow \max, \quad (18)$$

thus giving the random phase structure:

$$\Phi_j = \sum_{k=1}^n \arg \int d^3x \varphi_j^{*(k)}(t, \mathbf{r}) \varphi^{(k)}(\mathbf{r} - \mathbf{d}_j^{(k)}). \quad (19)$$

The random phase (19) can be used for simulating quantum computing via generating the following M random dichotomic functions:

$$f_s(\theta_s) = \text{sign} [\cos(\Phi_j + \theta_s)], \quad s = \overline{1, M}, \quad (20)$$

with θ_s being arbitrary fixed phases. Now recall that the quantum bit (qubit) is identified with the state vector

$$|\psi\rangle = \alpha|0\rangle + \beta|1\rangle,$$

corresponding to the superposition of two orthogonal states $|0\rangle$ and $|1\rangle$, as for instance, two polarizations of the photon, or two possible 1/2 - spin states. Using the well-known quantum expression for 1/2 - spin correlation in a singlet state of two particles (the latter being a typical entanglement state):

$$\mathbb{E}(\sigma_a \sigma_b) = -(\mathbf{a} \mathbf{b}), \quad |\mathbf{a}| = |\mathbf{b}| = 1, \quad (21)$$

one can compare it with the random phases correlation for the case of $n = 2$ particles [15]:

$$\mathbb{E}(f_1 f_2) = 1 - \frac{2}{\pi} |\Delta\theta|, \quad (22)$$

where $\Delta\theta = \theta_1 - \theta_2$. The similarity of these two functions (21) and (22) of the angular variable seems to be a good motivation for the M qubits simulation by the dichotomic random functions (20), popularized in the paper [15].

In conclusion we express the hope that the random solitons realization of the wave function could be effectively used for quantum computing simulation.

References

- [1] *de Broglie L.* Les Incertitudes d'Heisenberg et l'Interprétation Probabiliste de la Mécanique Ondulatoire. Paris: Gauthier-Villars, 1982; Moscow: Mir, 1986.- 342 p.
- [2] *Einstein A.* Collection of Scientific Works. Moscow: Nauka, vol. 4. 1967. - 600 p.
- [3] *Rybakov Yu. P.* The Bohm — Vigier Subquantum Fluctuations and Nonlinear Field Theory // Int. J. Theor. Phys., 1972, vol. 5, No. 2. Pp. 131 - 138.
- [4] *Rybakov Yu. P.* On the Causal Interpretation of Quantum Mechanics // Found. of Physics, 1974, vol. 4, No. 2. Pp. 149 - 161.
- [5] *Rybakov Yu. P.* La Théorie Statistique des Champs et la Mécanique Quantique // Ann. Fond. L. de Broglie, 1977, vol. 2, No. 3. Pp. 181 - 203.
- [6] *Rybakov Yu. P.* Self-Gravitating Solitons and Nonlinear - Resonance Quantization Mechanism // Bulletin of Peoples' Friendship University (PFU) of Russia. Physics. 1995, No. 3(1). Pp. 130 - 137.
- [7] *Rybakov Yu. P., Shachir M.* On Fresnel Diffraction of Solitons in the Singh Model // Izvestia VUZov. Physics, 1982, vol. 25, No. 1. Pp. 36 - 38.

- [8] *Rybakov Yu. P.* Gravitational Mechanism of Quantization and Solitons // Problems of Gravitation and Elementary Particles Theory. 1986, vol. 17. Pp. 161 - 171. Moscow: Energoatomizdat, 1986.
- [9] *Rybakov Yu. P., Kamalov T. F.* Stochastic Gravitational Fields and Quantum Correlations // Bulletin of PFU. Physics. 2002, No. 10(1). Pp. 5 - 7.
- [10] *Rybakov Yu. P., Saha B.* Soliton Model of Atom // Found. of Physics. 1995, vol. 25, No. 12. Pp. 1723 - 1731.
- [11] *Rybakov Yu. P., Saha B.* Interaction of a Charged 3D Soliton with a Coulomb Center // Phys. Lett., Ser. A, 1996, vol. 222, No. 1. Pp. 5 - 13.
- [12] *Taldykin A. T.* Vector Functions and Equations. Leningrad: LGU, 1977.- 352 p.
- [13] *Wiener N.* Differential Space // JMP, 1923, vol. 2, No. 3. Pp. 131 - 174.
- [14] *Wiener N.* Nonlinear Problems in Random Theory. N. Y.: MIT, 1958; Moscow: Inostr. Liter., 1961.- 160 p.
- [15] *Evdokimov N. V., Klyshko D. N., Komolov V. P., Yarochkin V. A.* Bell's Inequalities and EPR — Bohm Correlations: A Working Classical Radio Model // Progress in Physics (Russia), 1996, vol. 166, No. 1. Pp. 91 - 107.